


Genesee County
GENERAL TREATMENT
NAUSEA & VOMITING

Initial Date: 8/24/2012
Revised Date: 9/20/2019

Section 1-3

Nausea & Vomiting

1. Follow **General Pre-hospital Care Protocol**.

 2. For patients ≥ 40 kg that are not actively vomiting, administer Ondansetron (Zofran) 4mg ODT, per MCA selection.

a. **Contraindications:** Patients with Phenylketonuria (PKU)


ODT Ondansetron included?

YES NO

 3. For signs of dehydration, administer NS IV/IO fluid bolus up to 1 liter, wide open.


a. Pediatrics receive 20 ml/kg 

4. Hypotensive patients should receive additional IV/IO fluid boluses, as indicated by hemodynamic state. Continue IV/IO fluid bolus to a maximum of 2 liters.

a. Pediatrics repeat dose of 20 ml/kg 

 5. Administer Ondansetron (Zofran)

a. Adults 4mg IV/IM (if ODT not already administered or if patient vomited post ODT administration).

b. Pediatrics 0.1 mg/kg IV/IM, max dose of 4 mg 

 6. Repeat Ondansetron (Zofran) (**post-radio**)

a. Adults 4mg IV/IM (if ODT not already administered or if patient vomited post ODT administration).

b. Pediatrics 0.1 mg/kg IV/IM, max dose of 4 mg 